


# NCTN Lymphoma Trials Portfolio (Open as of 8/15/2025)

Each far right box includes the NCTN protocol number with a hyperlink to the associated ClinicalTrials.gov webpage. Click on it to view the protocol title and study information.



## NCTN Lymphoma Trials (Open as of 8/15/2025)

Protocol Number	Phase	Protocol Title
<b>A051901</b>	I	Phase I Trial of Methotrexate, Rituximab, Lenalidomide, and Nivolumab (Nivo-MR2) Induction Followed by Lenalidomide and Nivolumab Maintenance in Primary CNS Lymphoma
<b>A051902</b>	II	A Randomized Phase II Study of CHO(E)P vs CC-486-CHO(E)P vs Duvelisib-CHO(E)P in Previously Untreated CD30 Negative Peripheral T-Cell Lymphomas
<b>A052101</b>	III	A Randomized Phase 3 Trial of Continuous vs. Intermittent Maintenance Therapy with Zanubrutinib as Upfront Treatment in Older Patients with Mantle Cell Lymphoma
<b>AHOD2131</b>	III	A Randomized Phase 3 Interim Response Adapted Trial Comparing Standard Therapy with Immuno-oncology Therapy for Children and Adults with Newly Diagnosed Stage I and II Classic Hodgkin Lymphoma
<b>EA4231</b>	II	A Phase 2 Study of Venetoclax, Ibrutinib, Prednisone, Obinutuzumab, and Revlimid (VIPOR) in Relapsed or Refractory CD10-Negative Diffuse-Large B-Cell Lymphoma (DLBCL) and High-Grade B-Cell Lymphoma with MYC and BCL2 Rearrangements (HGBCL-DH-BCL2)
<b>EA4232</b>	III	A Randomized Phase III Study to Evaluate Benefits of Autologous Stem Cell Transplant in Patients with Peripheral T Cell Lymphoma That Achieved a First Complete Remission (CR1) Following Induction Therapy (PTCL-STAT)
<b>EA9213</b>	II	A Phase II Study of Daratumumab-Hyaluronidase for Chemotherapy-Relapsed/Refractory Minimal Residual Disease (MRD) in T Cell Acute Lymphoblastic Leukemia/Lymphoma (T-ALL/T-LBL)
<b>S1905</b>	II	A Phase I/II Study of AKR1C3-Activated Prodrug OBI-3424 (OBI-3424) In Patients with Relapsed/Refractory T-Cell Acute Lymphoblastic Leukemia (T-ALL)/T-Cell Lymphoblastic Lymphoma (T-LBL)
<b>S1918</b>	II/III	A Phase II/III Randomized Study of R-MiniCHOP with or Without CC-486 (Oral Azacitidine) in Participants Age 75 Years or Older with Newly Diagnosed Diffuse Large B Cell Lymphoma, Grade IIIB Follicular Lymphoma, Transformed Lymphoma, and High-Grade B-Cell Lymphomas with MYC AND BCL2 and/or BCL6 Rearrangements
<b>S1925</b>	III	Randomized, Phase III Study of Early Intervention with Venetoclax and Obinutuzumab Versus Delayed Therapy with Venetoclax and Obinutuzumab in Newly Diagnosed Asymptomatic High-Risk Patients with Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL): EVOLVE CLL/SLL Study
<b>S2005</b>	II	A Phase II Randomized Study of Ibrutinib and Rituximab with or Without Venetoclax in Previously Untreated Waldenström's Macroglobulinemia (WM) / Lymphoplasmacytic Lymphoma (LPL)
<b>S2114</b>	II	A Randomized Phase II Trial of Consolidation Therapy Following CD19 CAR T-Cell Treatment For Relapsed/Refractory Large B-Cell Lymphoma or Grade IIIB Follicular Lymphoma
<b>S2207</b>	II	Randomized Phase II Study of the Addition of Targeted Therapeutic Agents to Tafasitamab-Based Therapy in Non-Transplant-Eligible Patients with Relapsed/Refractory Large B-Cell Lymphoma
<b>S2308</b>	III	Randomized Phase III Study of Mosunetuzumab vs. Rituximab for Low Tumor Burden Follicular Lymphoma